

PSYCHOLOGICAL & SPECIALIST SERVICES

**Theory in the field: practice guidance for supporting  
young people experiencing suicidal distress and their  
carers**

---

PRESENTED BY KELSEY STALKER  
OCCUPATIONAL THERAPIST, LINKS TRAUMA HEALING SERVICE




# Acknowledgements

---


# Common misconceptions

Common beliefs	Facts
Only people who have depression die by suicide	Depression is a risk factor, however 'psychosocial risk factors' is the most commonly reported risk factor category in deaths by suicide
Suicidal thoughts are rare	22-38% of youth report having experienced suicidal thoughts
People who self-harm are attention seeking	A personal history of self harm is one of three most frequently occurring risk factors associated with death by suicide
All self-harming behaviour is suicidal behaviour	Self-harming behaviour can occur with or without the presence of suicidal thoughts/intention

# Psychosocial factors linked with suicide




# Further opportunity



# Survive then Thrive



# Survive



## Assess

Intent, plans and access to means  
Personal factors, context



## Act

Reduce access to means  
Increase psychosocial support  
Monitor




## Acute supports

Emergency Services (000)  
Hospital presentation  
NSW Mental Health Line (1800 011 511)  
Lifeline (13 11 14)

# Survive



**Conversations  
count**



**Collaborate**



**Language  
matters**



**Repair**



# Thrive




## Education

- Education maladjustment, difficulties with teachers /peers
- Education/literacy problems



## Culture

- Disruption of family
- Relationship problems: family
- Disappearance/death of family member
- Other problems (primary support group)



## Social supports

- Disruption of family
- Disappearance/death of family member
- Relationship problems: family, partner
- Difficulties with teachers/peers
- Other social problems



## Counselling/therapeutic supports

- Personal history of self harm
- History of sexual abuse



## Check in

- Longitudinal stressors

# Further supports/information

- **Lifeline**

- For individuals: <https://www.lifeline.org.au/get-help/i-m-feeling-suicidal/>
- For carers: <https://www.lifeline.org.au/get-help/i-m-worried-about-someone/>
- 13 11 14

- **Beyond Blue**

- For individuals: <https://www.beyondblue.org.au/mental-health/suicide-prevention/worried-about-someone-suicidal>
- For carers: <https://www.beyondblue.org.au/mental-health/suicide-prevention/worried-about-someone-suicidal>
- Various support options: [www.beyondblue.org.au](http://www.beyondblue.org.au)
- App (safety planning)

- **GP, Headspace**

- **Mental Health First Aid**

- Google search for services in your area

# References

- Australian Government. (2017). National Strategic Framework for Aboriginal and Torres Strait Islander Peoples' Mental Health and Social and Emotional Wellbeing 2017-2023.  
[https://www.niaa.gov.au/sites/default/files/publications/mhsewb-framework\\_0.pdf](https://www.niaa.gov.au/sites/default/files/publications/mhsewb-framework_0.pdf)
- Australian Institute of Health and Welfare. (2023, September 12). *Suicide & self-harm monitoring*.  
<https://www.aihw.gov.au/suicide-self-harm-monitoring/data/behaviours-risk-factors/psychosocial-risk-factors-suicide>
- Dazzi, T., Gribble, R., Wessely, S., & Fear, N. T. (2014). Does asking about suicide and related behaviours induce suicidal ideation? What is the evidence?. *Psychological medicine*, 44(16), 3361–3363.  
<https://doi.org/10.1017/S0033291714001299>
- Fortune, S., & Hetrick, S. (2022). Suicide risk assessments: Why are we still relying on these a decade after the evidence showed they perform poorly?. *The Australian and New Zealand journal of psychiatry*, 56(12), 1529–1534.  
<https://doi.org/10.1177/00048674221107316>
- Harmer, B., Lee, S., Duong, T. V. H., & Saadabadi, A. (2023). Suicidal Ideation. In *StatPearls*. StatPearls Publishing.
- Pisani, A. R., Murrie, D. C., & Silverman, M. M. (2016). Reformulating Suicide Risk Formulation: From Prediction to Prevention. *Academic psychiatry : the journal of the American Association of Directors of Psychiatric Residency Training and the Association for Academic Psychiatry*, 40(4), 623–629. <https://doi.org/10.1007/s40596-015-0434-6>
- Saini, P., Chantler, K., & Kapur, N. (2016). General practitioners' perspectives on primary care consultations for suicidal patients. *Health & social care in the community*, 24(3), 260–269. <https://doi.org/10.1111/hsc.12198>
- Suicide Prevention Australia (2023, September 14) *Stats and Facts*.  
<https://www.suicidepreventionaust.org/news/statsandfacts#1690952979591-101717b2-6937>
- Westerman, T. (2021) Culture-bound syndromes in Aboriginal Australian populations, *Clinical Psychologist*, 25:1, 19-35, DOI: [10.1080/13284207.2020.1843967](https://doi.org/10.1080/13284207.2020.1843967)